

TREASURY DEPARTMENT
UNITED STATES PUBLIC HEALTH SERVICE

HYGIENIC LABORATORY.—BULLETIN No. 93

APRIL, 1914

DIGEST OF COMMENTS
ON THE
PHARMACOPŒIA OF THE UNITED STATES
OF AMERICA
[EIGHTH DECENNIAL REVISION]
AND ON THE
NATIONAL FORMULARY
[THIRD EDITION]

FOR THE CALENDAR YEAR ENDING DECEMBER 31

1912

BY

MURRAY GALT MOTTER

AND

MARTIN I. WILBERT



WASHINGTON
GOVERNMENT PRINTING OFFICE
1914

CANNABIS INDICA.

Mitlacher, Wilhelm: Observations on the cultivation of *Cannabis sativa indica*.—Ztschr. allg. österr. Apoth.-Ver. 1912, v. 50, p. 377.

Borneman, John A.: Cannabis can be very readily cultivated.—Am. J. Pharm. 1912, v. 84, p. 553.

Lackey, R. H.: Bombay cannabis indica is practically unavailable at the present time and large quantities of other varieties of cannabis are being imported and apparently find a ready market.—Proc. Pennsylvania Pharm. Assoc. 1912, p. 61.

Pearson, W. A.: A Mexican and an American grown sample were found much below standard.—Proc. Pennsylvania Pharm. Assoc. 1912, p. 169.

Xrayser II.: Cannabis indica has fallen greatly into disuse in this country, and it matters little to us whether the drug is produced in Asia, Africa, or America.—Chem. & Drug. 1912, v. 81, p. 547.

Hamilton, H. C.: The pharmacopœial requirements for *C. sativa*, with comments on the need of further specifications.—J. Am. Pharm. Assoc. 1912, v. 1, p. 200-203.

Rusby, H. H.: Supplies of cannabis indica have continued scarce and prices high, which has stimulated the importation of large amounts of the drug which can not be regarded as the U. S. P. article.—J. Am. Pharm. Assoc. 1912, v. 1, p. 374.

Eckler and Miller: Commercial samples of American cannabis were found to vary widely in their activity. None were as active as good samples of the Indian drug.—Am. J. Pharm. 1912, v. 84, p. 488-495. See also Eighth Internat. Cong. Appl. Chem. 1912, v. 17, p. 23-30; and Pharm. Post, 1912, v. 45, p. 902.

Rippetoe and Minor: Six samples of cannabis indica contained from 14.20 to 14.44 per cent of ash.—Am. J. Pharm. 1912, v. 84, p. 437.

J. D. Riedel, A.-G.: The ash content of cannabis was found to vary from 6.0 to 7.1 per cent; amount of insoluble ash to 2.4 per cent.—Riedel's Berichte, 1912, p. 51.

Patch, E. L.: Samples of cannabis indica gave ether soluble resin, 12.1 per cent, 11.1 per cent.—J. Am. Pharm. Assoc. 1912, v. 1, p. 374.

Vanderkleed, Chas. E.: The assay of 4 samples of cannabis indica varied from 12.460 to 16.820 per cent of resin; all above standard.—Proc. Pennsylvania Pharm. Assoc. 1912, p. 179.

Mann, E. W.: A parcel of *Cannabis sativa*, grown in Greece, proved to contain a much higher proportion of resin than occurs in the Indian grown drug.—Ann. Rep. Southall Bros. & Barclay, 1912, 1913, p. 8.

Caesar & Loretz: Genuine Indian hemp was found to yield 17.53 per cent of extract while the African variety yielded 16.40 per cent.—Jahres-Ber. 1912, p. 62.

Stewart, F. E.: Fifteen samples of fluid extract of cannabis indica showed a variation of 40 to 150 per cent.—*J. Am. Pharm. Assoc.* 1912, v. 1, p. 1452.

Marshall and Wood: The value of the acetyl number in the standardization of preparations of Indian hemp.—*Brit. M. J.* 1912, v. 1, p. 1234.

Eckler, C. R.: Progress in methods of physiological testing; brief outline of the methods used in the standardization of the digitalis series, ergot and cannabis indica.—*Proc. Indiana Pharm. Assoc.* 1912, p. 65-70.

Houghton, E. M.: Report of the committee on physiological testing, on the physiological assay of pharmaceutical preparations of *C. sativa* by noting the nature of the intoxication in carefully selected dogs.—*J. Am. Pharm. Assoc.* 1912, v. 1, p. 1305.

Cushny, A. R.: Since the introduction of physiological assay by one firm they have had practically no complaints whatever as to the inefficacy of products of cannabis indica.—*Brit. M. J.* 1912, v. 2, p. 1558.

Moreau, B.: Brief note on Indian hemp. Apropos of the thesis of Bouquet.—*Bull. Sc. pharmacol.* 1912, v. 19, p. 599-601.

Book Review: An essay on hasheesh, including observations and experiments by Victor Robinson.—*Am. J. Pharm.* 1912, v. 84, p. 429.

Cook, N. M.: A specific use for cannabis indica is found in a condition simulating nervous breakdown and accompanied by irregular functions of the heart.—*Ellingwood's Therap.*, 1912, v. 6, p. 171-172.